



Soil Carbon Sequestration Impacts on Global Climate Change and Food Security

R. Lal, *et al.*
Science **304**, 1623 (2004);
DOI: 10.1126/science.1097396

The following resources related to this article are available online at www.sciencemag.org (this information is current as of February 6, 2008):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/304/5677/1623>

Supporting Online Material can be found at:

<http://www.sciencemag.org/cgi/content/full/304/5677/1623/DC1>

A list of selected additional articles on the Science Web sites related to this article can be found at:

<http://www.sciencemag.org/cgi/content/full/304/5677/1623#related-content>

This article cites 26 articles, 6 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/304/5677/1623#otherarticles>

This article has been cited by 107 article(s) on the ISI Web of Science.

This article has been cited by 10 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/cgi/content/full/304/5677/1623#otherarticles>

This article appears in the following subject collections:

Ecology

<http://www.sciencemag.org/cgi/collection/ecology>

Information about obtaining reprints of this article or about obtaining permission to reproduce this article in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

The carbon sink capacity of the world's agricultural and degraded soils is 50 to 66% of the historic carbon loss of 42 to 78 gigatons of carbon. The rate of soil organic carbon sequestration with adoption of recommended technologies depends on soil texture and structure, rainfall, temperature, farming system, and soil management. Strategies to increase the soil carbon pool include soil restoration and woodland regeneration, no-till farming, cover crops, nutrient management, manuring and sludge application, improved grazing, water conservation and harvesting, efficient irrigation, agroforestry practices, and growing energy crops on spare lands. An increase of 1 ton of soil carbon pool of degraded cropland soils may increase crop yield by 20 to 40 kilograms per hectare (kg/ha) for wheat, 10 to 20 kg/ha for maize, and 0.5 to 1 kg/ha for cowpeas. As well as enhancing food security, carbon sequestration has the potential to offset fossil-fuel emissions by 0.4 to 1.2 gigatons of carbon per year, or 5 to 15% of the global fossil-fuel emissions.